

SYMPLECTIC AND POISSON GEOMETRY ON b -MANIFOLDS

VICTOR GUILLEMIN, EVA MIRANDA, AND ANA RITA PIRES

ABSTRACT. Let M^{2n} be a Poisson manifold with Poisson bivector field Π . We say that M is b -Poisson if the map $\Pi^n : M \rightarrow \Lambda^{2n}(TM)$ intersects the zero section transversally on a codimension one submanifold $Z \subset M$. This paper will be a systematic investigation of such Poisson manifolds. In particular, we will study in detail the structure of (M, Π) in the neighbourhood of Z and using symplectic techniques define topological invariants which determine the structure up to isomorphism. We also investigate a variant of de Rham theory for these manifolds and its connection with Poisson cohomology.

1. INTRODUCTION

In her 2002 paper [R], Radko gave a classification of stable Poisson structures on a compact 2-manifold M , where stable for a Poisson bivector field Π means that the map

$$\Pi : M \rightarrow \Lambda^2(TM)$$

is transverse to the zero section. In this paper we describe some partial generalizations of this result to higher dimensions, with the stability condition replaced by a more complicated condition which we call b -Poisson: the map

$$\Pi^n = \Pi \wedge \dots \wedge \Pi : M \rightarrow \Lambda^{2n}(TM)$$

must be transverse to the zero section of $\Lambda^{2n}(TM)$. We denote by Z the hypersurface in M where this map is zero. Then, Π restricted to Z defines a regular Poisson structure Π_Z on Z with codimension-one symplectic leaves.

When generalizing the results in [R] to n dimensions, one is confronted with the question of whether or not a such a Poisson structure on Z can be extended to a b -Poisson structure on a neighbourhood of Z in M . We use modular geometry techniques of Weinstein to give necessary and sufficient conditions on Π_Z for this to be the case. Then, we address the more complicated issue of how many extensions exist up to Poisson isomorphism and show that these are classified by the elements of $H_{\text{Poisson}}^1(Z)$. To prove

Date: March 7, 2013.

Eva Miranda is partially supported by the DGICYT/FEDER project MTM2009-07594: Estructuras Geometricas: Deformaciones, Singularidades y Geometria Integral. Ana Rita Pires has the partial support of an AMS-Simons Travel Grant.

this we give an alternative definition of b -Poisson: we show following Melrose and Nest-Tsygan that a b -Poisson manifold can be regarded dually as a b -symplectic manifold. More explicitly, we define a b -manifold to be a pair (M, Z) , where M is a manifold and Z a (not necessarily connected) hypersurface in M . We define a b -category in which the objects are such pairs and the morphisms are maps

$$f : (M_1, Z_1) \rightarrow (M_2, Z_2)$$

where f is transverse to Z_2 and $Z_1 = f^{-1}(Z_2)$; we call these b -maps. If one defines the b -tangent bundle bTM and b -cotangent bundle ${}^bT^*M$ for (M, Z) following [Me]¹, these have nice functorial properties with respect to b -maps. We then define a b -symplectic form to be a nondegenerate closed b -two form ω , i.e., a section of $\Lambda^2({}^bT^*M)$ which is nondegenerate at all $p \in Z$ and show that $\omega|_Z$ defines a Poisson vector field v on (Z, Π_Z) and hence a cohomology class in the quotient

$$H_{\text{Poisson}}^1(Z) = \frac{\text{Poisson vector fields}}{\text{Hamiltonian vector fields}}.$$

Our proof that the converse assertion is true, i.e., that $[v]$ determines up to isomorphism the b -Poisson structure on M (locally in a neighbourhood of Z), is then reduced to a standard Moser type argument: if ω_0 and ω_1 are two b -symplectic forms on a tubular neighbourhood \mathcal{U} of Z in M and $\omega_0|_Z = \omega_1|_Z$, then $\omega_0 - \omega_1 = d\mu$ for some b -one form $\mu \in {}^b\Omega^1(\mathcal{U})$, and one gets a diffeomorphism on \mathcal{U} mapping ω_0 to ω_1 by integrating the b -vector field v_t defined by

$$\iota_{v_t}\omega_t = \mu, \text{ where } \omega_t = (1 - t)\omega_0 + t\omega_1.$$

We also use this Moser technique to prove that for ω_0 and ω_1 globally defined on a compact M , ω_0 and ω_1 are globally b -symplectomorphic if the usual global Moser conditions are satisfied, i.e., if ω_t is b -symplectic and $[\omega_t] \in {}^bH^2(M, \mathbb{R})$ is independent of t , in addition to $\omega_t|_Z$ being independent of t . We will show that in dimension two this argument gives an alternative b -symplectic proof of Radko's theorem.

The main part of this paper (sections 2 to 8) is a more detailed account of these results. In sections 2 and 3 we review and introduce a number of basic definitions in b -geometry, from b -manifolds to b -differential forms. In section 4 we examine the notion of b -symplectic from the Poisson-geometer's perspective, prove that b -symplectic and b -Poisson are equivalent notions and investigate some consequences of this fact: these include the relation

¹As is described in section 2, we have borrowed the name of b -manifolds from the b -calculus developed by Melrose [Me]. Another possible notation is inspired from Algebraic Geometry and the approach of Goto [Go] and, more recently, by Gualtieri and Li in [GL] in the holomorphic case. Our approach to b -geometry in this paper is close to but not exactly the same as that of Melrose, where a b -manifold is a manifold with boundary and $Z = \partial M$.

between Weinstein's splitting theorem and a b -version of Moser's theorem, as well as the role of Weinstein's modular vector field in this theory. In section 5, we discuss Mazzeo-Melrose's b -analogue of the de Rham theorem, obtain some cohomology results for b -symplectic manifolds and compare it to two versions of Poisson cohomology. In section 6 we discuss Darboux and Moser theorems for b -symplectic manifolds and use these b -methods to prove Radko's theorem, in section 7 we revisit the modular invariants introduced in [GMP], and in section 8 show that these b -methods enable us to generalize Radko's theorem to arbitrary dimension, by addressing the extension problem that we alluded to above: does a regular Poisson structure on Z extend to a b -symplectic structure on M , and if so in how many different ways?

All these results seem to put the b -symplectic category closer to the symplectic world than to the usually cumbersome Poisson world².

Section 9 of this paper is devoted to issues that we will investigate in more detail in the future. In section 9 we show that b -symplectic manifolds are integrable, i.e., that the Lie algebroid structure on M defined by Π can always be integrated to a symplectic groupoid structure. This turns out to be an easy corollary of a much more general result [CF03] of Crainic-Fernandes, in a future paper we will describe in more detail examples of symplectic groupoids whose existence is a consequence of this result. We also give an explicit model for the symplectic groupoid integrating the exceptional hypersurface Z .

Finally, in section 10, we discuss completely integrable systems on b -symplectic manifolds, i.e., systems f_1, \dots, f_n of b -Poisson commuting functions which are almost everywhere functionally independent. In dimension two, f can be chosen to be nonsingular at points of Z and in fact to be a defining function for Z . In dimension four, f_1 can be chosen generically to be a defining function for Z and f_2 to have standard elliptic and hyperbolic singularities on the (2-dimensional) symplectic leaves of Z . Analogous results can be obtained for higher dimensions. In a future paper we plan to study integrable systems and Hamiltonian actions in the b -symplectic context.

Acknowledgements: We have benefited from many interesting, useful and lively discussions with the following people: Marius Crainic, Rui Loja Fernandes, Marco Gualtieri, David Martinez Torres, Ryszard Nest and Francisco Presas. These discussions have enriched and improved this paper.

²In this sense, it is interesting to point out that an h -principle holds also for b -symplectic manifolds as it has been recently proved by Freijlich [F]

2. WHAT DOES b STAND FOR?

We recall the notions of b -tangent and b -cotangent bundles introduced by Richard Melrose in [Me] as a framework to study differential calculus and differential operators on manifolds with boundary, and further studied by Ryszard Nest and Boris Tsygan in [NT] in the context of formal deformations of symplectic manifolds with boundary.

Let $(M, \partial M)$ be a manifold with boundary and consider $p \in \partial M$. A neighbourhood of p is diffeomorphic to the upper half n -space

$$H_+^n = \{(x_1, \dots, x_n) | x_1 \geq 0\},$$

and the tangent space at p is generated by $x_1 \frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}, \dots, \frac{\partial}{\partial x_n}$. Intuitively this means that vector fields on $(M, \partial M)$ are vector fields on M which are tangent to ∂M along the boundary. More formally, we can use sheaves to define the tangent and cotangent bundles as follows.

Let $U \subset M$ be an open set. The set of vector fields on U which are tangent to ∂M is denoted $\Gamma(U, {}^b\mathcal{T})$, and by varying the open set U we obtain a sheaf on M . Observe that for any point $p \in \partial(H_+^n)$, the tangent bundle at p is generated by $T_p(H_+^n) = \langle x_1 \frac{\partial}{\partial x_1}_p, \frac{\partial}{\partial x_2}_p, \dots, \frac{\partial}{\partial x_n}_p \rangle$. Therefore, for open sets $U \subset M$ diffeomorphic to H_+^n , we have $\Gamma(U, {}^b\mathcal{T})$ freely generated over $C^\infty(M)$ by sections $x_1 \frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}, \dots, \frac{\partial}{\partial x_n}$.

Definition 1. The b -tangent bundle ${}^bT(M)$ of a manifold with boundary $(M, \partial M)$ is the vector bundle associated to the locally free sheaf $\Gamma(U, {}^b\mathcal{T})$. The b -cotangent bundle ${}^bT^*(M)$ is the dual bundle to ${}^bT(M)$.

Note that the b -cotangent bundle ${}^bT^*(M)$ is locally generated by the sections: $\frac{dx_1}{x_1}, dx_2, \dots, dx_n$.

The manifolds which we study in this paper are not manifolds with boundary, but rather manifolds with a distinguished hypersurface. As we will see in section 3, the appropriate notions of tangent and cotangent bundles are basically the ones above³. For this reason, we adopt the notation bTM and ${}^bT^*M$ and furthermore name our objects of study “ b -manifolds”.

Some results proved for manifolds with boundary hold under appropriate translation for our b -manifolds, for example the Darboux theorem for closed nondegenerate 2 forms [NT] and the Mazzeo-Melrose splitting of the cohomology groups of $(M, \partial M)$ in terms of the cohomology of M and of ∂M [Me], as we will see in the next sections.

³This was pointed out in [CW] §17.4 in the context of giving an example of a Lie algebroid associated to the bundle of vectors tangent to M which are furthermore tangent to a fixed hypersurface of M .

3. DIFFERENTIAL FORMS ON b -MANIFOLDS

In this section we introduce b -manifolds and b -maps, which are respectively the objects and morphisms of the b -category, revisit the notions of b -tangent and b -cotangent bundles. We then define differential b -forms and construct a b -de Rham cohomology theory.

3.1. The b -tangent and b -cotangent bundles. We begin with some definitions.

Definition 2. A **b -manifold** is a pair (M, Z) of an oriented manifold M and an oriented hypersurface $Z \subset M$. A **b -map** is a map $f : (M_1, Z_1) \rightarrow (M_2, Z_2)$ transverse to Z_2 and such that $f^{-1}(Z_2) = Z_1$. The **b -category** is the category whose objects are b -manifolds and morphisms are b -maps.

Definition 3. A **b -vector field** on M is a vector field which is tangent to Z at every point $p \in Z$.

These vector fields form a Lie subalgebra of the algebra of all vector fields on M . Moreover, they also form a projective module over the ring $C^\infty(M)$, and hence are sections of a vector bundle on M . We call this vector bundle the **b -tangent bundle** and denote it bTM .

If v is a b -vector field, then the restriction $v|_Z$ is everywhere tangent to Z , and hence defines a vector field v_Z on Z . Thus we have a map $\Gamma({}^bTM|_Z) \rightarrow \Gamma(TZ)$ and since this map is a morphism of $C^\infty(Z)$ -modules, it is induced by a vector bundle morphism

$$(1) \quad {}^bTM|_Z \rightarrow TZ.$$

Proposition 4. *The kernel of the map (1) is a line bundle \mathbb{L}_Z with a canonical nonvanishing section.*

Proof. Since Z is oriented, there exists a defining function for Z , i.e. a b -map $f : (M, Z) \rightarrow (\mathbb{R}, 0)$. Let v be a vector field on M with $df_p(v_p) = 1$ for all $p \in Z$. Then $w = fv|_Z$ is a nonvanishing section of \mathbb{L}_Z , and it is easily checked that its definition doesn't depend on the choice of the function f or the vector field v . \square

We call this nonvanishing section w of \mathbb{L}_Z the **normal b -vector field** of the b -manifold (M, Z) .

Note that at points $p \in M \setminus Z$, the b -tangent space coincides with the usual tangent space ${}^bT_pM = T_pM$, whereas at points $p \in Z$, there is a surjective map

$${}^bT_pM \rightarrow T_pZ$$

with kernel spanned by w_p .

We define the ***b*-cotangent bundle** of M to be the vector bundle ${}^bT^*M$ dual to bTM . Then, at points $p \in M \setminus Z$, the *b*-cotangent space coincides with the usual cotangent space: ${}^bT_p^*M = T_p^*M$. At points $p \in Z$, there is an embedding

$$T_p^*Z \rightarrow {}^bT_p^*M$$

whose image is $\{l \in {}^bT_p^*M \mid l(w_p) = 0\}$.

Given a defining function f for Z , let $\mu \in \Omega^1(M \setminus Z)$ be the one-form $\frac{df}{f}$. If v is a *b*-vector field then the pairing $\langle v, \mu \rangle \in C^\infty(M \setminus Z)$ extends smoothly over Z and hence μ itself extends smoothly over Z as a section of ${}^bT^*M$. Moreover, $\mu_p(w_p) = 1$ for $p \in Z$, so given f we get a splitting

$$(2) \quad {}^bT_p^*M = T_p^*Z + \text{span}\{\mu_p\}.$$

For ease of notation, we will write $\mu_p = \frac{df_p}{f}$, even though the expression on the right hand side is not well-defined for $p \in Z$.

3.2. The *b*-de Rham complex. For each k , let ${}^b\Omega^k(M)$ denote the space of ***b*-de Rham k -forms**, i.e., sections of the vector bundle $\Lambda^k({}^bT^*M)$. The usual space of de Rham k -forms sits inside this space in a somewhat non-trivial way: given $\mu \in \Omega^k(M)$, we interpret it as a section of $\Lambda^k({}^bT^*M)$ by the convention

$$\begin{aligned} \mu_p &\in \Lambda^k(T_p^*M) = \Lambda^k({}^bT_p^*M) \text{ at } p \in M \setminus Z \\ \mu_p &= (i^*\mu)_p \in \Lambda^k(T_p^*Z) \subset \Lambda^k({}^bT_p^*M) \text{ at } p \in Z \end{aligned}$$

where $i : Z \hookrightarrow M$ is the inclusion map.

With these conventions it is easy to see that every *b*-de Rham k -form can be written as

$$(3) \quad \omega = \alpha \wedge \frac{df}{f} + \beta, \text{ with } \alpha \in \Omega^{k-1}(M) \text{ and } \beta \in \Omega^k(M).$$

Moreover, while α and β are not unique, we claim:

Proposition 5. *At $p \in Z$, α_p and β_p are unique.*

Proof. This follows from the direct sum decomposition (2) and the fact that at $p \in Z$, α_p and β_p have to be interpreted as elements of $\Lambda^{k-1}(T_p^*Z)$ and $\Lambda^k(T_p^*Z)$. \square

The decomposition (3) enables us to extend the exterior d operator to ${}^b\Omega(M)$ by setting

$$d\omega = d\alpha \wedge \frac{df}{f} + d\beta.$$

The right hand side is well defined and agrees with the usual exterior d operator on $M \setminus Z$ and also extends smoothly over M as a section of $\Lambda^{k+1}({}^bT^*M)$.

Note that $d^2 = 0$, which allows us to form a complex of b -forms, the b -de Rham complex:

$$0 \rightarrow {}^b\Omega^0(M) \xrightarrow{d} {}^b\Omega^1(M) \xrightarrow{d} {}^b\Omega^2(M) \xrightarrow{d} \dots \rightarrow 0$$

Also note that if $\omega = \alpha \wedge \frac{df}{f} + \beta \in {}^b\Omega^k(M)$ is closed, it is not necessarily true that α and β are closed, take for example $\beta = 0$ and $\alpha = \mu \wedge df$ with μ chosen so that α is not closed.

In general, b -de Rham forms “explode” at Z . Those that vanish at Z are in fact honest de Rham forms:

Proposition 6. *If $\omega \in {}^b\Omega^k(M)$ is such that $\omega|_Z = 0$, then $\omega \in \Omega^k(M)$.*

Proof. Given $\omega = \alpha \wedge \frac{df}{f} + \beta \in {}^b\Omega^k(M)$, the condition $\omega|_Z = 0$ implies that $\alpha|_Z = 0$ and $\beta|_Z = 0$. If $\alpha|_Z = 0$ then $\alpha_1 := \frac{\alpha}{f}$ is in ${}^b\Omega^{k-1}(M)$. Thus $\omega = \alpha_1 \wedge df + \beta$ is in $\Omega^k(M)$. \square

Remark 7. Even though b -forms “explode” at Z , it is possible to integrate compactly supported b -forms of top degree over M : for $\omega \in {}^b\Omega^d(M)$ compactly supported we define the integral of ω over M to be

$$\int_M \omega = \lim_{\varepsilon \rightarrow 0} \int_{|f| > \varepsilon} \omega,$$

where f is a defining function for Z . This limit exists and is independent of the choice of f , the proof is similar to that of Theorem 2 in [R].

4. b -SYMPLECTIC MANIFOLDS AND b -POISSON MANIFOLDS

4.1. b -symplectic manifolds. In this section we introduce the notion of symplectic for the b -category, and observe that b -symplectic manifolds are also Poisson manifolds. A simple example is the b -manifold (M, Z) where $M = \mathbb{R}^{2n}$ with coordinates $x_1, y_1, \dots, x_n, y_n$ and Z is the hyperplane $x_1 = 0$. Consider the closed b -form

$$\omega = x_1 \wedge \frac{dy_1}{y_1} + \sum_{i=2}^n dx_i \wedge dy_i.$$

This form is non degenerate in the sense that ω^n is a well defined, nonvanishing b -form. That this example is the local prototype of all b -symplectic manifolds is the content of the b -Darboux theorem (Theorem 32).

Dualizing ω we obtain the bivector field

$$\Pi = y_1 \frac{\partial}{\partial x_1} \wedge \frac{\partial}{\partial y_1} + \sum_{i=2}^n \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_i},$$

which is a Poisson structure on M since $[\Pi, \Pi] = 0$. The symplectic foliation for this Poisson structure contains two open leaves — the upper and lower half spaces given by $y_1 > 0$ and $y_1 < 0$ —, and the union of the remaining leaves is the hyperplane Z , where Π^n vanishes — these leaves are $(2n - 2)$ -dimensional planes through the different levels of x_1 .

Definition 8. Let (M, Z) be a $2n$ -dimensional b -manifold and $\omega \in {}^b\Omega^2(M)$ a closed b -form. We say that ω is **b -symplectic** if ω_p is of maximal rank as an element of $\Lambda^2({}^bT_p^*M)$ for all $p \in M$.

Example 9. Analogous to what happens in the symplectic case, the b -cotangent bundle of a b -manifold (M, Z) is a b -symplectic manifold. If y_i are local coordinates for the manifold M on a neighbourhood of a point in Z , with Z defined locally by $y_1 = 0$, and x_i are the fiber coordinates on ${}^bT^*M$, then the canonical one-form is given in these coordinates by

$$x_1 \frac{dy_1}{y_1} + \sum_{i=2}^n x_i dy_i,$$

and its exterior derivative

$$dx_1 \wedge \frac{dy_1}{y_1} + \sum_{i=2}^n dx_i \wedge dy_i$$

is a b -symplectic form on ${}^bT^*M$.

As seen in (3), fixing a defining function f for Z we can decompose the symplectic form as

$$(4) \quad \omega = \alpha \wedge \frac{df}{f} + \beta, \text{ with } \alpha \in \Omega^1(M) \text{ and } \beta \in \Omega^2(M).$$

Proposition 10. Let $\tilde{\alpha} = i^*\alpha$ and $\tilde{\beta} = i^*\beta$, where $i : Z \hookrightarrow M$ is the inclusion. Then

1. The forms $\tilde{\alpha}$ and $\tilde{\beta}$ are closed.
2. The form $\tilde{\alpha}$ is nowhere vanishing and intrinsically defined in the sense that it does not depend on the splitting (4). In particular, the codimension-one foliation of Z defined by $\tilde{\alpha}$ is intrinsically defined.
3. For each leaf $L \xrightarrow{i_L} Z$ of this foliation, the form $i_L^*\tilde{\beta}$ is intrinsically defined, and is a symplectic form on L .

Proof. 1. From (4) we get $d\alpha \wedge \frac{df}{f} + d\beta = 0$, and by Proposition 5, the forms $d\alpha_p$ and $d\beta_p$ are zero as elements of $\Lambda^2(T_p^*Z)$ and $\Lambda^3(T_p^*Z)$ for all $p \in Z$, i.e., $di^*\alpha$ and $di^*\beta$ are zero.

2. As seen in (2), the cotangent space ${}^bT_p^*M$ for $p \in Z$ is the direct sum $T_p^*Z + \text{span} \left\{ \frac{df_p}{f} \right\}$. Thus, if we had $\tilde{\alpha}_p = 0$, we would have $\omega_p = \tilde{\beta}_p \in \Lambda^2(T_p^*Z)$, implying $\text{rank}(\omega_p) < 2n$, so $\tilde{\alpha}$ is nonvanishing. Replacing f by

another defining function g , we have $f = gh$, with h nonvanishing on Z . Near Z we have $\frac{df}{f} = \frac{dg}{g} + d(\log |h|)$ and (4) becomes

$$\omega = \alpha \wedge \frac{dg}{g} + \beta_1,$$

where $\beta_1 = \beta + \alpha \wedge d(\log |h|)$. Then, for $\tilde{\beta}_1 = \tilde{\beta} + \tilde{\alpha} \wedge d(\log |h|)|_Z$,

$$(5) \quad \omega|_Z = \tilde{\alpha} \wedge \frac{dg}{g}|_Z + \tilde{\beta}_1.$$

3. For a leaf L of the foliation defined by $\tilde{\alpha}$, by (5) we have $i_L^* \tilde{\beta}_1 = i_L^* \tilde{\beta}$. If $i_L^* \tilde{\beta}$ were of rank smaller than $(2n-2)$ at some $p \in Z$, then $\omega_p = \tilde{\alpha}_p \wedge \frac{df_p}{f}$ would be of rank smaller than n as an element of ${}^b\Lambda^2(T_p^*M)$.

□

Note that by Proposition 5, the two-form $\tilde{\beta}$ is a symplectic invariant of the pair (ω, f) . Moreover, (5) describes how $\tilde{\beta}$ depends on the choice of the defining function.

4.2. b -Poisson manifolds. We now look at the Poisson counterparts of b -symplectic manifolds, and will prove in the next section that these two notions are equivalent. This notion corresponds to non-degenerate Poisson structures on b -manifolds which for short we call b -Poisson structures. One could consider other Poisson structures on pairs (M, Z) but the ones described below are the object of this paper.

Definition 11. Let (M^{2n}, Π) be an oriented Poisson manifold such that the map

$$p \in M \mapsto (\Pi(p))^n \in \Lambda^{2n}(TM)$$

is transverse to the zero section, then $Z = \{p \in M | (\Pi(p))^n = 0\}$ is a hypersurface and we say that Π is a **b -Poisson structure** on (M, Z) and (M, Z) is a **b -Poisson manifold**.

In this subsection we give some examples (compact and otherwise) of b -Poisson structures. In section 8 we will find a procedure to construct examples of b -Poisson manifolds from regular Poisson manifolds with certain vanishing obstruction classes.

Example 12. The Lie algebra of the affine group of dimension 2 is a model for noncommutative Lie algebras in dimension 2, its algebra structure is given by $[e_1, e_2] = e_2$, where e_1, e_2 form a basis. We can naturally write this (bilinear) Lie algebra structure as the Poisson structure

$$\Pi = y \frac{\partial}{\partial x} \wedge \frac{\partial}{\partial y},$$

which is dual to the b -symplectic form $\omega = dx \wedge \frac{dy}{y}$. In this example the critical hypersurface Z is the y -axis, it is the union of symplectic leaves of dimension 0 (points on the line), and the upper and lower half-planes are symplectic open leaves of dimension 2.

Example 13. On the sphere \mathbb{S}^2 with the usual coordinates (h, θ) , the Poisson structure $\Pi = h \frac{\partial}{\partial h} \wedge \frac{\partial}{\partial \theta}$ vanishes transversally along the equator $Z = \{h = 0\}$ and hence is a b -Poisson structure on (\mathbb{S}^2, Z) . We can define a b -Poisson structure on any orientable surface by choosing a curve and a defining function for it.

In dimension 2, these structures were studied and classified by Radko in [R] under the name of *topologically stable Poisson structures*. In this case, Z is a union of smooth curves and each point in these curves is a symplectic leaf of Z . In [R], Radko proves that the following ingredients give a complete classification of b -Poisson manifolds of dimension 2:

- The set of curves $\gamma_1, \dots, \gamma_n$ along which the Poisson structure vanishes;
- The periods along the curves $\gamma_1, \dots, \gamma_n$ of a modular vector field⁴ on M associated to the volume form ω_Π , the two-form dual to Π , on the complement of Z ;
- The regularized Liouville volume of (M, Π) , which is a correction along Z of the natural volume associated to the Poisson structure, necessary because the original volume form ω_Π associated to the Poisson structure Π blows up at Z . This regularized Liouville volume is the integral $\int_M \omega_\Pi$ as defined in Remark 7.

The following classification holds:

Theorem 14 (Radko). *The set of curves, modular periods and regularized Liouville volume completely determines, up to Poisson diffeomorphisms, the b -Poisson structure on a compact surface M .*

In the paper [R], the author then uses this classification to explicitly compute the Poisson cohomology of the manifold in terms of the modular vector field.

Example 15. So far our examples have been of dimension 2, but product structures allow us to get to higher dimensions: let (R, π_R) be a Radko compact surface and (S, π_S) be a compact symplectic surface, then $(R \times S, \pi_R + \pi_S)$ is a b -Poisson manifold of dimension 4. Furthermore, perturbations of product structures can be used to obtain non-product ones. For instance, take \mathbb{S}^2 with b -Poisson structure $\Pi_1 = h \frac{\partial}{\partial h} \wedge \frac{\partial}{\partial \theta}$ and the symplectic torus \mathbb{T}^2

⁴For a precise definition of modular vector field see section 4.4.

with dual Poisson structure $\Pi_2 = \frac{\partial}{\partial \theta_1} \wedge \frac{\partial}{\partial \theta_2}$. Then

$$\hat{\Pi} = h \frac{\partial}{\partial h} \wedge \left(\frac{\partial}{\partial \theta} + \frac{\partial}{\partial \theta_1} \right) + \Pi_2.$$

is a b -Poisson structure on $\mathbb{S}^2 \times \mathbb{T}^2$, but not the product one described above.

We will see in Section 6 that using Moser path methods in the b -context, we can control perturbations that produce equivalent Poisson structures.

Example 16. Let (N^{2n+1}, π) be a regular corank-1 Poisson manifold, X be a Poisson vector field and $f : \mathbb{S}^1 \rightarrow \mathbb{R}$ a smooth function. The bivector field

$$\Pi = f(\theta) \frac{\partial}{\partial \theta} \wedge X + \pi$$

is a b -Poisson structure on $\mathbb{S}^1 \times N$ if the function f vanishes linearly and the vector field X is transverse to the symplectic leaves of N (this condition is necessary to guarantee transversality). If that is so, the critical hypersurface consists of the union of as many copies of N as zeros of f .

In this example N^{2n+1} is the exceptional hypersurface of the b -Poisson manifold and has an induced Poisson structure which is regular of corank one. It is a general fact that the exceptional hypersurface of a b -Poisson manifold naturally inherits a corank-one Poisson structure. This example provides the semilocal model for a b -Poisson structure in a neighbourhood of the critical hypersurface Z .

4.3. b -Poisson equals b -symplectic. In this section, we will show the following:

Proposition 17. *A two-form ω on a b -manifold (M, Z) is b -symplectic if and only if its dual bivector field Π is a b -Poisson structure.*

We begin by recalling Weinstein's splitting theorem, which we will then apply to the particular case of b -Poisson manifolds:

Theorem 18 (Weinstein). *Let (M^m, Π) be a Poisson manifold of rank $2k$ at a point $p \in M$. Then there exists a neighborhood and a local coordinate system $(x_1, y_1, \dots, x_k, y_k, z_1, \dots, z_{m-2k})$ centered at p for which the Poisson structure can be written as*

$$(6) \quad \Pi = \sum_{i=1}^k \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_i} + \sum_{i,j=1}^{m-2k} f_{ij}(z) \frac{\partial}{\partial z_i} \wedge \frac{\partial}{\partial z_j},$$

where f_{ij} are functions which depend only on the variables (z_1, \dots, z_{m-2k}) and which vanish at the origin.

Let Π be a b -Poisson structure on (M, Z) , and Ω and Ξ a volume form on M and its dual $2n$ -vector field, respectively. Then, $\Pi^n = f\Xi$ for some $f : M \rightarrow \mathbb{R}$ which vanishes on Z . Since the $2n$ -vector field Π^n doesn't

vanish identically, the generic rank of the Poisson structure is $2n$, and it is less than $2n$ on Z . This implies that the two-form ω_Π dual to Π is a smooth symplectic form on $M \setminus Z$. Because Π^n intersects the zero section of $\bigwedge^n(TM)$ transversally, 0 is a regular value of f and so $Z = f^{-1}(0)$ must be a codimension-one submanifold of M , a union of hypersurfaces. Furthermore, we can assume that in a neighbourhood of a point in Z , the function f is simply the coordinate function z_1 , with $z_1 = 0$ locally defining the hypersurface. When restricted to Z , the Poisson structure defines a symplectic foliation of codimension one. To prove this, use Theorem 18 and observe that Π^n vanishing transversally at Z implies that the transverse Poisson manifolds at points of Z must be of dimension two, so Z is the union of symplectic leaves of corank 2 in M . This defines a codimension-one foliation of Z by symplectic leaves.

Summarizing, a b -Poisson manifold is a Poisson manifold (M^{2n}, Π) for which the $2n$ -vector field Π^n vanishes linearly along a union of smooth hypersurfaces Z and such that the Poisson structure Π defines a symplectic structure ω_Π on $M \setminus Z$ and when restricted to Z gives a symplectic foliation of codimension one in Z . In particular, the rank maximality of a b -symplectic form implies that its dual bivector field must be a b -Poisson structure; this proves one of the directions of Proposition 17.

Consider now the particular case of a 2-dimensional b -Poisson manifold (M, Z) . The Poisson bivector field Π vanishes linearly at Z , and the dual two-form will be given locally by $\omega_\Pi = \frac{1}{z_1 f(z_1, z_2)} dz_1 \wedge dz_2$, with $f(z_1, z_2)$ nonvanishing on Z , which is given locally by $z_1 = 0$. The diffeomorphism ϕ given by the change of coordinates $z = z_1$ and $t = \int f(z_1, z_2) dz_2$ satisfies $\phi^*(\omega_\Pi) = \frac{1}{z} dz \wedge dt$ (here we give the explicit diffeomorphism, but the existence of such a diffeomorphism derives simply from the fact that Π^n intersects the zero section $\bigwedge^n(TM)$ transversally and uses the regular value theorem).

Combining this result with Theorem 18, we obtain a local normal form result resembling the Darboux theorem, which will appear again as Theorem 32 and that we will reprove then using Moser path methods.

Proposition 19. *Let (M, Z) be a b -Poisson manifold, with Poisson bivector field Π and dual two-form ω_Π . Then, on a neighbourhood of a point $p \in Z$, there exist coordinates $(x_1, y_1, \dots, x_{n-1}, y_{n-1}, z, t)$ centered at p such that*

$$\omega_\Pi = \sum_{i=1}^{n-1} dx_i \wedge dy_i + \frac{1}{z} dz \wedge dt.$$

Remark 20. In other words, we can find a splitting such that

$$\omega_\Pi = \omega_L + (\Pi^T)^\sharp$$

where ω_L is the symplectic form on the symplectic leaf through the point $p \in Z$ and $(\Pi^T)^\sharp$ is the dual to a b -Poisson structure on a 2-dimensional

manifold. In particular, we have obtained a linearization result for bivector fields associated to b -manifolds.

Because being symplectic is a local property, Proposition 19 implies that a b -Poisson manifold is b -symplectic, the other direction of Proposition 17. From now on, we will refer to these manifolds as b -symplectic manifolds.

4.4. Modular vector fields of b -symplectic manifolds. A modular vector field on a Poisson manifold measures how far Hamiltonian vector fields are from preserving a given volume form. A simple example is that of a symplectic manifold endowed with the volume form that is the top power of its symplectic form: the modular vector field will be zero because this volume form is invariant under the flow of any Hamiltonian vector field. In this section we study modular vector fields of b -symplectic manifolds.

We follow Weinstein [We97] for the description of modular vector fields of Poisson manifold; a complete presentation of these can also be found in [K]. Some results about modular vector fields for regular corank one Poisson manifolds, as is the case of the exceptional hypersurface Z of a b -symplectic manifold (M, Z) , can be found in [GMP].

Definition 21. Let (M, Π) be a Poisson manifold and Ω a volume form on it, and denote by u_f the Hamiltonian vector field associated to a smooth function f on M . The **modular vector field** X_Π^Ω (or simply X^Ω if the Poisson structure is fixed and hence implicit) is the derivation given by the mapping

$$f \mapsto \frac{\mathcal{L}_{u_f} \Omega}{\Omega}.$$

Let (M, Z) be a b -symplectic manifold and consider the local coordinates given by Proposition 19 in a neighbourhood of a point $p \in Z$. The b -symplectic form ω can be written as

$$\omega = \sum_{i=1}^{n-1} dx_i \wedge dy_i + \frac{1}{z} dz \wedge dt,$$

and consider also the volume form

$$\Omega = dx_1 \wedge dy_1 \wedge \cdots \wedge dx_{n-1} \wedge dy_{n-1} \wedge dz \wedge dt.$$

Working in these local coordinates we see that the modular vector field associated to the volume form Ω and the Poisson structure dual to the b -symplectic form is given by

$$X^\Omega = \frac{\partial}{\partial t}.$$

As a consequence we have:

Proposition 22. *The modular vector field of a b -symplectic manifold (M, Z) is tangent to Z and transverse to the symplectic leaves inside Z , independently of the volume form considered on M .*

Proof. At each point $p \in Z$, working in the local coordinates mentioned above, the modular vector field with respect to the volume form Ω above is

$$X^\Omega = \frac{\partial}{\partial t}.$$

This vector field is tangent to Z , which is given locally by $z = 0$, and transverse to the symplectic foliation inside Z , because the leaves of that foliation are locally just the different levels of the coordinate function t .

If we consider another volume form $\Omega' = H\Omega$, where $H \in C^\infty(M)$ is nonvanishing, the modular vector field becomes

$$X^{\Omega'} = \frac{\partial}{\partial t} + u_{\log(|H|)},$$

it differs from the previous one by a hamiltonian vector field⁵. Hamiltonian vector fields are tangent to the symplectic leaves of M , and in particular, to all the $(2n - 2)$ -dimensional leaves whose union is Z and hence to Z itself. Therefore, the new modular vector field $X^{\Omega'}$ will still be tangent to Z and transverse to the symplectic leaves in it. \square

Using these local coordinates we also see that $\tilde{\alpha}(v_{\text{mod}}|_Z) = 1$, independently of choice of modular vector field v_{mod} .

5. COHOMOLOGY THEORIES FOR b -MANIFOLDS

In this section we explore some cohomology theories for b -manifolds, and the relationships between them. For a b -manifold (M, Z) , we can talk about the usual cohomology theories for the underlying manifold M , such as de Rham cohomology and Poisson cohomology, which correspond respectively to de Rham forms and to multivector fields, but we can also use the notions of b -forms and b -multivector fields, and study the corresponding cohomology theories.

5.1. De Rham cohomology and b -cohomology. We begin by proving a Mazzeo-Melrose theorem for b -manifolds (see [Me], §2.16, page 88 for the original version) and then as a direct application obtain some results about low degree b -cohomology for b -symplectic manifolds.

⁵The **modular class** of a Poisson manifold is the class of a modular vector field in the first Poisson cohomology group, it depends on the Poisson structure but not on the volume form: a change in the volume form changes the modular vector field by a hamiltonian vector field.

Let (M, Z) be a b -manifold with $Z \xrightarrow{i} M$ compact.

Theorem 23. [*b -Mazzeo-Melrose theorem*] *The b -cohomology groups of M are computable by*

$${}^bH^*(M) \cong H^*(M) \oplus H^{*-1}(Z).$$

Proof. Let $f : (M, Z) \rightarrow (\mathbb{R}, 0)$ be a defining function for Z . Then, every $\omega \in {}^b\Omega^k(M)$ can be written as $\omega = \alpha \wedge \frac{df}{f} + \beta$, with $\alpha, \beta \in \Omega^*(M)$. Moreover, in b -de Rham theory, $\tilde{\alpha} := i^*\alpha$, where $i : Z \hookrightarrow M$ is the inclusion, is intrinsically defined independent of the choice of f , so we have a canonical short exact sequence of de Rham complexes

$$0 \rightarrow \Omega^k(M) \rightarrow {}^b\Omega^k(M) \rightarrow \Omega^{k-1}(Z) \rightarrow 0$$

which induces a long exact sequence in cohomology

$$\dots \rightarrow H^k(M) \xrightarrow{i} {}^bH^k(M) \xrightarrow{j} H^{k-1}(Z) \xrightarrow{\delta} H^{k-1}(M) \rightarrow \dots$$

This sequence splits into short exact sequences because that the map j is surjective: let $\mathcal{U} \cong Z \times (-\varepsilon, \varepsilon)$ be a collar neighbourhood of Z in M , change f so that $f \equiv 1$ on the complement of $Z \times (-\frac{\varepsilon}{2}, \frac{\varepsilon}{2})$, and let $p : Z \times (-\frac{\varepsilon}{2}, \frac{\varepsilon}{2}) \rightarrow Z$ be the projection. Then, for any closed $\tilde{\alpha} \in \Omega^{k-1}(Z)$, the form $\omega = p^*\tilde{\alpha} \wedge \frac{df}{f} \in {}^b\Omega^k(M)$, and $j[\omega] = [\tilde{\alpha}]$. \square

Observe that if M is compact then the theorem above says that for cohomology of top dimension we have

$${}^bH^d(M) = H^d(M) \oplus (\oplus_i H^{d-1}(Z_i))$$

where the Z_1, Z_2, \dots, Z_r are the connected components of Z , and hence $\dim({}^bH^d(M)) = r + 1$.

The symplectic form on a compact symplectic manifold defines a non-vanishing second cohomology class on the manifold, yielding a non-trivial second cohomology group. For b -symplectic manifolds, the analogue involves the second b -cohomology group:

Proposition 24. *For a compact b -symplectic manifold (M, Z) we have $H^1(Z) \neq \{0\}$ and consequently ${}^bH^2(M) \neq \{0\}$.*

Proof. The submanifold Z , being the vanishing set of Π^n (the top power of the bivector field dual to the b -symplectic form), is closed and since M is compact, Z is compact as well. Let α be a one-form Z that defines the corank-1 regular foliation induced by Π on Z : the form α is nowhere vanishing and $i_L^*\alpha = 0$ for all leaves $L \xrightarrow{i_L} Z$. If we had $H^1(Z) = 0$ then α would be exact: $\alpha = dg$ for some function $g \in C^\infty(Z)$. By compactness of Z , the function g has maximum and minimum points, at which $\alpha = dg$ would then necessarily vanish.

Then, by Theorem 23 we have

$${}^bH^2(M) \cong H^2(M) \oplus H^1(Z) \neq \{0\}.$$

□

Similarly, we have:

Proposition 25. *For a compact b -symplectic manifold (M, Z) we have $H^2(Z) \neq \{0\}$ and consequently ${}^bH^3(M) \neq \{0\}$.*

Proof. Suppose that $H^2(Z) = 0$. Then $d\tilde{\beta} = 0$ on Z implies that $\tilde{\beta} = d\mu$ for some one-form μ defined on Z . Since $\tilde{\beta}^n \wedge \tilde{\alpha}$ is a volume form on Z (see [GMP]), we have

$$0 \neq \text{vol}(Z) = \int_Z \tilde{\beta}^n \wedge \tilde{\alpha} = \int_Z (d\mu)^n \wedge \tilde{\alpha} = \int_{\partial Z} \mu \wedge (d\mu)^{n-1} \wedge \tilde{\alpha} = 0.$$

Thus $H^2(Z)$ must be non-trivial and Theorem 23 gives us ${}^bH^3(M) \neq \{0\}$. □

5.2. Poisson cohomology and b -Poisson cohomology. In this section we study the relationship between b -cohomology and Poisson cohomology, obtain a Mazzeo-Melrose type result for Poisson cohomology and as a direct application rederive the computation of Poisson cohomology for two-dimensional b -manifolds obtained in [R].

For a general Poisson manifold (M, Π) , the Poisson structure Π induces a differential operator $d_\Pi = [\Pi, \cdot]$ on the graded algebra of multivector fields on M by extending the Lie bracket to multivectorfields. The cohomology of the complex of multivector fields $\Lambda^*(M)$

$$\dots \longrightarrow \Lambda^{k-1}(M) \xrightarrow{d_\Pi} \Lambda^k(M) \xrightarrow{d_\Pi} \Lambda^{k+1}(M) \longrightarrow \dots$$

is the Poisson cohomology $H_\Pi^*(M)$ of M associated with the Poisson structure Π .

Let ${}^b\Lambda^k(M)$ denote the space of **b -multivector fields**, i.e., sections of the vector bundle $\Lambda^k({}^bTM)$. Then the operator $d_\Pi = [\Pi, \cdot]$ is a differential on the subalgebra of b -multivector fields on M . The **b -Poisson cohomology** ${}^bH_\Pi^*(M)$ associated to the b -Poisson structure Π on M is the cohomology of the complex

$$\dots \longrightarrow {}^b\Lambda^{k-1}(M) \xrightarrow{d_\Pi} {}^b\Lambda^k(M) \xrightarrow{d_\Pi} {}^b\Lambda^{k+1}(M) \longrightarrow \dots$$

Explicit computations of Poisson cohomology are close to impossible in the general Poisson case. A simple example in which Poisson cohomology can be computed is the case of symplectic manifolds, for which the Poisson cohomology is isomorphic to de Rham cohomology, this is because non-degeneracy of the symplectic form allows us to define a bundle isomorphism

between T^*M and TM . Similarly, in the b -symplectic case, non-degeneracy of the b -symplectic form gives a bundle isomorphism between ${}^bT^*M$ and bTM , which translates to an isomorphism between b -de Rham cohomology and b -Poisson cohomology.

Theorem 26. *Let (M, Z) be a b -symplectic manifold, and Π the corresponding b -Poisson structure. Then, the b -Poisson cohomology ${}^bH_{\Pi}^*(M)$ is isomorphic to the b -de Rham cohomology ${}^bH^*(M)$.*

Proof. We define the operator $\natural : {}^bT^*M \rightarrow {}^bTM$ such that for $\alpha, \beta \in {}^b\Omega^1(M)$ we have

$$\langle \alpha \wedge \beta, \Pi \rangle = \langle \beta, \natural(\alpha) \rangle.$$

By taking exterior powers of \natural we obtain for each k a homomorphism between $\Lambda^k({}^bT^*M)$ and $\Lambda^k({}^bTM)$, and hence also between ${}^b\Omega^k(M)$ and ${}^b\Lambda^k(M)$. Because the Poisson structure Π is non-degenerate in this b -context, this homomorphism is an isomorphism, which we will also denote by \natural .

The classical formula for Lichnerowicz complexes

$$\natural(d\eta) = -[\Pi, \natural(\eta)] = -d_{\Pi}(\natural(\eta))$$

guarantees that the following diagram commutes

$$\begin{array}{ccccccc} \dots & \longrightarrow & {}^b\Lambda^{k-1}(M) & \xrightarrow{d_{\Pi}} & {}^b\Lambda^k(M) & \xrightarrow{d_{\Pi}} & {}^b\Lambda^{k+1}(M) \longrightarrow \dots \\ & & \uparrow \natural & & \uparrow \natural & & \uparrow \natural \\ \dots & \longrightarrow & {}^b\Omega^{k-1}(M) & \xrightarrow{d} & {}^b\Omega^k(M) & \xrightarrow{d} & {}^b\Omega^{k+1}(M) \longrightarrow \dots \end{array}$$

thus providing the desired isomorphism between the cohomologies of the two complexes. \square

We are interested not only in the b -Poisson cohomology of a b -symplectic manifold (M, Z) , but also in its honest Poisson cohomology. We will see that the difference between the bundles $\Lambda^*(M)$ and ${}^b\Lambda^*(M)$ translates into a contribution of the Poisson cohomology of Z , producing a formula reminiscent of that of Mazzeo-Melrose's.

Theorem 27. *Let (M, Z) be a compact b -symplectic manifold, Π the corresponding b -Poisson structure, and Π_Z the restriction of Π to Z . Then the Poisson cohomology groups of M are computable by*

$$H_{\Pi}^k(M) \cong {}^bH^k(M) \oplus H_{\Pi_Z}^{k-1}(Z).$$

Proof. The set of b -multivector fields on (M, Z) is naturally included in the set of multivector fields on M . Fixing a defining function $f : (M, Z) \rightarrow (\mathbb{R}, 0)$ for Z , we can associate to a k -vector field on M , a $(k-1)$ -vector field on Z , by contracting with df and restricting to Z . Note that contracting a

b -multivector field with df and restricting to Z yields the 0 multivector field on Z , so we have a (non-canonical) short exact sequence of complexes:

$$0 \rightarrow {}^b\Lambda^k(M) \rightarrow \Lambda^k(M) \rightarrow \Lambda^{k-1}(Z) \rightarrow 0,$$

which induces a long exact sequence in cohomology

$$\dots \rightarrow {}^bH_{\Pi}^k(M) \xrightarrow{i} H_{\Pi}^k(M) \xrightarrow{p} H_{\Pi_Z}^{k-1}(Z) \xrightarrow{\delta} {}^bH_{\Pi}^{k-1}(M) \rightarrow \dots$$

This sequence splits into short exact sequences because the map p is surjective: let A be a $(k-1)$ -vector field on Z and \hat{A} an extension of it to M . Then $\frac{\partial}{\partial f} \wedge \hat{A}$ is a k -vector field on M whose image under p is A . Furthermore, if A is a cochain $\Lambda^{k-1}(Z)$, then it is also a cochain in the complex $\Lambda^k(M)$ because

$$d_{\Pi}(\frac{\partial}{\partial f} \wedge \hat{A}) = [\Pi, \frac{\partial}{\partial f} \wedge \hat{A}] = [\Pi, \frac{\partial}{\partial f}] \wedge \hat{A} - \frac{\partial}{\partial f} \wedge [\Pi, \hat{A}] = 0 - \frac{\partial}{\partial f} \wedge d_{\Pi_Z}(A).$$

Thus, we have $H_{\Pi}^k(M) \cong {}^bH_{\Pi}^k(M) \oplus H_{\Pi_Z}^{k-1}(Z)$ and the desired result is obtained by using Theorem 26 on the ${}^bH_{\Pi}^k(M)$ term. \square

Remark 28. When the symplectic foliation of Z has compact leaves, the Poisson structure of Z is that of a symplectic mapping torus as proved in [GMP] (see also section 7 in this paper). One can then use the results of Vaisman [V90, V94] to describe the Poisson cohomology of Z in terms of its de Rham cohomology, the set of smooth functions on a submanifold $N \subset Z$ transversal to the symplectic foliation, and the set of vector fields tangent to N . These computations can also be recovered from [X] because the exceptional hypersurface is integrable as proved in [GMP] (see section 7).

In the two-dimensional case, we can use Theorem 27 to reprove a result of Radko's [R].

Corollary 29 (Radko). *Let (M, Z) be a compact connected two-dimensional b -symplectic manifold, where M is of genus g and Z a union of n curves on M . Then the Poisson cohomology of M is given by*

$$\begin{aligned} H_{\Pi}^0(M) &= \mathbb{R} \\ H_{\Pi}^1(M) &= \mathbb{R}^{n+2g} \\ H_{\Pi}^2(M) &= \mathbb{R}^{n+1}. \end{aligned}$$

Proof. Because Z is a union of curves, all Poisson cohomology groups $H_{\Pi_Z}^k(Z)$ vanish. Then, by Theorems 27 and 23, we have

$$H_{\Pi}^k(M) \cong {}^bH^k(M) \cong H^k(M) \oplus H^{k-1}(Z).$$

The result is then immediate from the fact that the nonzero cohomology groups of an oriented compact connected surface M of genus g are $H^0(M) =$

$H^2(M) = \mathbb{R}$ and $H^1(M) = \mathbb{R}^{2g}$, and those of a union Z of n curves are $H^0(Z) = H^1(Z) = \mathbb{R}^n$. \square

6. NORMAL FORMS

With the necessary tools and notions now in place, we can prove b -analogues of standard symplectic geometry theorems.

6.1. Relative Moser theorem for b -symplectic manifolds.

Theorem 30. *Let ω_0 and ω_1 be two b -symplectic forms on (M, Z) . If $\omega_0|_Z = \omega_1|_Z$, then there exist neighbourhoods $\mathcal{U}_0, \mathcal{U}_1$ of Z in M and a diffeomorphism $\gamma : \mathcal{U}_0 \rightarrow \mathcal{U}_1$ such that $\gamma|_Z = \text{id}_Z$ and $\gamma^*\omega_1 = \omega_0$.*

Recall that for $p \in Z$, one should interpret $\omega|_p$ as sitting in $\Lambda^2(T_p^*Z)$.

Proof. (using Moser trick) Let $\omega_t = (1 - t)\omega_0 + t\omega_1$. We will prove that there exists a neighbourhood \mathcal{U} of Z in M and an isotopy $\gamma_t : \mathcal{U} \rightarrow M$, with $0 \leq t \leq 1$ such that $\gamma_t|_Z = \text{id}_Z$ and

$$(7) \quad \gamma_t^*\omega_t = \omega_0.$$

If such a γ_t is to exist, by differentiating (7) we will get

$$(8) \quad \mathcal{L}_{v_t}\omega_t = \omega_0 - \omega_1,$$

where $v_t = \frac{d\gamma_t}{dt} \circ \gamma_t^{-1}$. Note that since $\gamma_t|_Z = \text{id}_Z$, we would have $v_t|_Z = 0$, so v_t would be a b -vector field vanishing on Z .

Because $(\omega_0 - \omega_1)|_Z = 0$, by Proposition 6 the b -form $(\omega_0 - \omega_1)$ is also an honest de Rham form, and since it is closed, by the Poincaré lemma there exists a one-form $\mu \in \Omega^1(M)$ such that $(\omega_0 - \omega_1) = d(f\mu)$ on a neighbourhood of Z , where $f : (M, Z) \rightarrow (\mathbb{R}, 0)$ is a defining function for Z . Then, (8) becomes

$$(9) \quad \iota_{v_t}\omega_t = f\mu,$$

which can be solved for v_t in a small enough neighbourhood \mathcal{U} of Z where ω_t is b -symplectic. Moreover, since the right hand side of (9) vanishes at points of Z , the b -vector field v_t thus defined does too.

We can get a suitable γ_t by integrating v_t , and the vector field vanishing on Z implies that $\gamma_t|_Z = \text{id}_Z$ as desired. Now set $\gamma := \gamma_1$ and the open sets $\mathcal{U}_0 := \mathcal{U}$ and $\mathcal{U}_1 := \gamma_1(\mathcal{U})$. \square

An alternative statement of Theorem 30 is the following:

Theorem 31. *Let ω_0 and ω_1 be two b -symplectic forms on (M, Z) . If they induce on Z the same restriction of the Poisson structure and their modular vector fields differ on Z by a hamiltonian vector field, then there exist neighbourhoods $\mathcal{U}_0, \mathcal{U}_1$ of Z in M and a diffeomorphism $\gamma : \mathcal{U}_0 \rightarrow \mathcal{U}_1$ such that $\gamma|_Z = \text{id}_Z$ and $\gamma^*\omega_1 = \omega_0$.*

Proof. It suffices to show that $\omega_0|_Z = \omega_1|_Z$, then apply Theorem 30.

Fix a defining function f for Z and write $\omega_j = \alpha_j \wedge \frac{df}{f} + \beta_j$, and also $\tilde{\alpha}_j = i^*\alpha_j$ and $\tilde{\beta}_j = i^*\beta_j$, with $j = 0, 1$ and $i : Z \hookrightarrow M$ the inclusion. What we want to show is that $\tilde{\alpha}_0 = \tilde{\alpha}_1$ and $\tilde{\beta}_0 = \tilde{\beta}_1$. Note that $v_{\text{mod } j}$ being the modular vector field implies that $\tilde{\alpha}_j(v_{\text{mod } j}|_Z) = 1$ and $\iota_{v_{\text{mod } j}|_Z}\tilde{\beta}_j = 0$.

From $\tilde{\alpha}_j(v_{\text{mod } j}|_Z) = 1$ and $(v_{\text{mod } 0}|_Z - v_{\text{mod } 1}|_Z)$ being a hamiltonian vector field on Z , we get $\tilde{\alpha}_0 = \tilde{\alpha}_1$.

Because ω_0 and ω_1 induce the same restriction of Poisson structure on the hypersurface Z , we have $i_L^*\tilde{\beta}_0 = i_L^*\tilde{\beta}_1$ for any symplectic leaf L , with $i_L^* : L \hookrightarrow Z$ the inclusion. This, together $\iota_{v_{\text{mod } j}|_Z}\tilde{\beta}_j = 0$, gives $\tilde{\beta}_0 = \tilde{\beta}_1$. \square

6.2. Darboux theorem for b -symplectic manifolds. As in the symplectic case, the relative Moser theorem can be used to prove a local canonical form result for b -symplectic forms, an analogue of the classical Darboux theorem.

Theorem 32. [b -Darboux theorem] *Let ω be a b -symplectic form on (M, Z) , and p a point in Z . Then we can find a coordinate system $(\mathcal{U}, x_1, y_1, \dots, x_n, y_n)$ centered at p such that on \mathcal{U} the hypersurface Z is locally defined by $y_1 = 0$ and*

$$\omega = dx_1 \wedge \frac{dy_1}{y_1} + \sum_{i=2}^n dx_i \wedge dy_i.$$

Proof. Let $\omega = \alpha \wedge \frac{df}{f} + \beta$, and $\tilde{\alpha} = i^*\alpha$ and $\tilde{\beta} = i^*\beta$, where $i : Z \hookrightarrow M$ is the inclusion. As seen in Proposition 10, for all $p \in Z$ we have $\tilde{\alpha}_p$ nonvanishing, $\tilde{\alpha}_p \wedge \tilde{\beta}_p \neq 0$ and $\tilde{\beta}_p \in \Lambda^2(T_p^*Z)$ of rank $n - 1$. Hence, by the standard Darboux theorem for closed two-forms of constant rank, there exists a coordinate system $(\mathcal{V}, x_1, x_2, y_2, \dots, x_n, y_n)$ on Z centered at p such that in these coordinates $\tilde{\alpha} = dx_1$ and $\tilde{\beta} = \sum_{i=2}^n dx_i \wedge y_i$. Then, if we let $y_1 = f$ we have on a neighbourhood \mathcal{U} of p in M whose intersection with Z is contained in \mathcal{V} ,

$$\omega|_Z = (dx_1 \wedge \frac{dy_1}{y_1} + \sum_{i=2}^n dx_i \wedge dy_i)|_Z.$$

The desired result now follows from Theorem 30. \square

6.3. Global Moser theorem for b -symplectic manifolds. When M is compact, we obtain a global result:

Theorem 33. [*b -Moser theorem, v.1*] *Suppose that M is compact and let ω_0 and ω_1 be two b -symplectic forms on (M, Z) . Suppose that ω_t , for $0 \leq t \leq 1$, is a smooth family of b -symplectic forms on (M, Z) joining ω_0 and ω_1 and such that the b -cohomology class $[\omega_t]$ and the restriction $\omega_t|_Z$ are independent of t . Then, there exists a family of diffeomorphisms $\gamma_t : M \rightarrow M$, for $0 \leq t \leq 1$ such that $\gamma_t|_Z = \text{id}_Z$ and $\gamma_t^* \omega_t = \omega_0$.*

Proof. As in the proof of Theorem 30, the existence of the desired isotopy γ_t relies on the existence of a smooth family of b -vector fields v_t such that

$$(10) \quad \mathcal{L}_{v_t} \omega_t = \frac{d\omega_t}{dt}.$$

Integrating this family of b -vector fields will then yield a suitable γ_t .

Because $[\omega_t]$ is independent of t , we have $[\frac{d\omega_t}{dt}] = \frac{d[\omega_t]}{dt} = 0$, and so there exists a family $\mu_t \in {}^b\Omega^1(M)$ such that $\frac{d\omega_t}{dt} = \mu_t$. Furthermore, because $\omega_t|_Z$ is also independent of t , we have $\mu_t|_Z = 0$, and thus by Proposition 6, this is in fact a smooth⁶ family of honest de Rham forms $\mu_t \in \Omega^1(M)$ which vanish on Z .

Equation 10 becomes

$$\iota_{v_t} \omega_t = \mu_t$$

which can be solved for v_t on M because the ω_t are b -symplectic. Furthermore, because the μ_t are a smooth family and vanish at Z , the v_t will be a smooth family b -vector fields, as desired. \square

Using the argument in the proof of Theorem 31 we can replace, in the statement of Theorem 33, the condition of $\omega_t|_Z$ being independent of t by all the ω_t inducing on Z the same restriction of the Poisson structure and the same restriction of the modular vector field.

Theorem 34. [*b -Moser theorem, v.2*] *Suppose that M is compact and let ω_0 and ω_1 be two b -symplectic forms on (M, Z) . Suppose that ω_t , for $0 \leq t \leq 1$, is a smooth family of b -symplectic forms on (M, Z) joining ω_0 and ω_1 and such that the b -cohomology class $[\omega_t]$, the restriction to Z of the Poisson structure induced by ω_t , and the restriction to Z of the modular vector field (up to addition of hamiltonian vector fields) induced by ω_t are all independent of t . Then, there exists a family of diffeomorphisms $\gamma_t : M \rightarrow M$, for $0 \leq t \leq 1$ such that $\gamma_t|_Z = \text{id}_Z$ and $\gamma_t^* \omega_t = \omega_0$.*

⁶The proof of smoothness is as described in the proof of Theorem 7.3 in [Ca].

6.4. Revisiting Radko's classification theorem. In the classical symplectic world, the Moser theorem applied to the two-dimensional case tells us that surfaces with the same symplectic volume must be symplectomorphic. The b -version of the Moser theorem, Theorem 33 (or its slightly different formulation Theorem 34), yields as its two-dimensional instance the classification of Radko surfaces.

Let M be a compact two-dimensional manifold and Π_0 and Π_1 be two Radko surface structures on it which vanish along the same set of curves $\gamma_1, \dots, \gamma_n$, induce the same modular periods along these curves and have the same regularized Liouville volume. Let us prove, as a corollary of Theorem 33 that the respective dual b -symplectic forms ω_0 and ω_1 on $(M, Z = \bigcup_{i=1}^n \gamma_i)$ are b -symplectomorphic.

For $j = 0, 1$, write $\omega_j = \alpha_j \wedge \frac{df_j}{f_j} + \beta_j$. We can assume that f_0 and f_1 have the same sign, since $\frac{d(-f_i)}{-f_i} = \frac{df_i}{f_i}$, and then using a diffeomorphism, that $f_0 = f_1 = f$. Also, since M is 2-dimensional, β_j must be of the form $\beta_j = \mu_j \wedge df$, for some one-form μ_j , and so by renaming $\alpha_j + f\mu_j$ to α_j , which note doesn't change its restriction to Z , we can write

$$\omega_j = \alpha_j \wedge \frac{df}{f}.$$

Define the convex combination $\omega_t = (1-t)\omega_0 + t\omega_1 = \alpha_t \wedge \frac{df}{f}$, where $\alpha_t = (1-t)\alpha_0 + t\alpha_1$. We want to show that the restriction $\omega_t|_Z$ and the b -cohomology class $[\omega_t]$ are both independent of t , and that for all $0 \leq t \leq 1$, the form ω_t is b -symplectic.

Firstly, the modular vector field is unique up to adding hamiltonian vector fields, which are tangent to the symplectic leaves. In dimension two, because the leaves contained in Z are points, the modular vector field will be unique. Furthermore, a vector field on a closed simple curve is uniquely determined (up to diffeomorphism) by its period, and thus the periods of the modular vector field along the curves $\gamma_1, \dots, \gamma_n$ completely determine the restriction of the modular vector field to Z . Thus, $v_{\text{mod } 0}|_Z = v_{\text{mod } 1}|_Z$. Then for $\tilde{\alpha}_j = i^* \alpha_j$, with $i : Z \hookrightarrow M$, which is characterized by $\tilde{\alpha}_j(v_{\text{mod } j}|_Z) = 1$, we have $\tilde{\alpha}_0 = \tilde{\alpha}_1$, and thus $\omega_0|_Z = \omega_1|_Z$ and so $\omega_t|_Z$ is independent of t .

Secondly, because $\omega_0|_Z = \omega_1|_Z$ and Proposition 6, the two-form $(\omega_0 - \omega_1)$ is an honest de Rham form. The regularized Liouville volumes associated with ω_0 and ω_1 being equal means that

$$\int_M (\omega_0 - \omega_1) = 0$$

both as a b -integral (see Remark 7) and, since it is an honest de Rham form that we are integrating, as an honest integral. Thus, the (honest de Rham)

cohomology class of $[\omega_0 - \omega_1]$ is zero, and thus so is its b -cohomology class. Therefore, $[\omega_t] = [\omega_0] = [\omega_1] \in {}^b H^2(M)$.

Lastly, we must show that ω_t is a b -symplectic form for all t . Away from Z , the forms ω_0 and ω_1 are honest area forms inducing the same orientation (the orientation is induced by the modular vector field's restriction to Z), and on Z , $\tilde{\alpha}_t = \tilde{\alpha}_0 = \tilde{\alpha}_1$ does not vanish, and thus $\omega_t = \alpha_t \wedge \frac{df}{f}$ is nondegenerate and hence b -symplectic.

7. INVARIANTS ASSOCIATED TO THE EXCEPTIONAL HYPERSURFACE OF A b -SYMPLECTIC MANIFOLD

Given a b -symplectic structure, the exceptional hypersurface is endowed with a Poisson structure which is regular of corank one. In this section we study and we completely characterize the foliation invariants of the codimension one symplectic foliation on Z . These invariants of Z were introduced in [GMP] in the context of codimension one symplectic foliation. The definition of these invariants does not require the existence of a b -symplectic structure on (M, Z) or even that of a manifold M of which Z is a hypersurface. The first invariant requires a transversally orientable codimension-one regular foliation on Z , the second one a Poisson structure on Z which induces a symplectic foliation with those characteristics. When Z is the exceptional hypersurface of a b -symplectic manifold (M, Z) , we get for free such a foliation and Poisson structure on Z .

7.1. The defining one-form of a foliation and the first obstruction class. Let Z be an odd-dimensional manifold, and \mathcal{F} a transversally orientable codimension-one foliation on Z .

Definition 35. A form $\alpha \in \Omega^1(Z)$ is a **defining one-form** of the foliation \mathcal{F} if it is nowhere vanishing and $i_L^* \alpha = 0$ for all leaves $L \xrightarrow{i_L} Z$.

In particular, when the foliation \mathcal{F} on Z is the symplectic foliation induced on Z by a b -symplectic structure on (M, Z) , a defining one-form can be chosen such that $\alpha(v_{mod}|_Z) = 1$. With this extra condition, the defining one-form is unique even when we consider a different volume form on M : this causes the modular vector field v_{mod} to change by a hamiltonian vector field, which is tangent to the leaves of \mathcal{F} . Also in this particular case, the modular one-form α will necessarily be closed:

Proposition 36. *If (M, Z) is a b -symplectic manifold then a defining one-form α of the symplectic foliation on Z that satisfies $\alpha(v_{mod}|_Z) = 1$ is necessarily closed.*

Proof. The flow of v_{mod} preserves the foliation \mathcal{F} , so it also preserves corresponding defining one-form α : $\mathcal{L}_{v_{mod}} \alpha = 0$. Applying Cartan's formula

we have

$$0 = \mathcal{L}_{v_{mod}|_Z} \alpha = d\iota_{v_{mod}|_Z} \alpha + \iota_{v_{mod}|_Z} d\alpha.$$

The first summand vanishes because $\alpha(v_{mod}|_Z) = 1$, and so the second summand vanishes as well: $(d\alpha)(v_{mod}|_Z, -) = 0$.

Since for a point $p \in Z$ we have the decomposition

$$T_p Z = \text{span}(v_{mod}|_Z) \oplus T_p L,$$

where L is the corank 2 symplectic leaf through p , it only remains to check that $d\alpha(v_1, v_2) = 0$ for $v_1, v_2 \in T_p L$. But $di_L^* \alpha = i_L^* d\alpha$ and therefore $d\alpha(v_1, v_2) = 0$. \square

We follow [GMP] for the definition the first obstruction class, an invariant related to the defining one-form of a foliation \mathcal{F} on Z . Consider the short exact sequence of complexes

$$0 \longrightarrow \alpha \wedge \Omega(Z) \xrightarrow{i} \Omega(Z) \xrightarrow{j} \Omega(Z)/\alpha \wedge \Omega(Z) \longrightarrow 0.$$

Because $i_L^* d\alpha = 0$ for all $L \in \mathcal{F}$ and α is a defining one-form of the foliation \mathcal{F} , we have

$$d\alpha = \beta \wedge \alpha \text{ for some } \beta \in \Omega^1(Z).$$

Furthermore, because $0 = d(d\alpha) = d\beta \wedge \alpha - \beta \wedge d\alpha = d\beta \wedge \alpha$, the two-form $d\beta$ is in the subcomplex $\alpha \wedge \Omega(Z)$, and so $d(j\beta) = 0$.

Definition 37. The **first obstruction class** of the foliation \mathcal{F} is

$$c_{\mathcal{F}} = [j\beta] \in H^1(\Omega(Z)/\alpha \wedge \Omega(Z)).$$

Note that since a different defining one-form will be $\alpha' = f\alpha$ with f a nonvanishing function, the complexes $\alpha \wedge \Omega(Z)$ and $\Omega(Z)/\alpha \wedge \Omega(Z)$ do not depend on the choice of α . Furthermore, one can prove that the class $c_{\mathcal{F}}$ is independent of the choice of α and that:

Theorem 38. [GMP] *The first obstruction class vanishes, $c_{\mathcal{F}} = 0$, if and only if one can choose the defining one-form α of \mathcal{F} to be closed.*

Remark 39. As it was proved in [GMP], manifolds with vanishing first obstruction class are unimodular (its modular class vanishes).

In particular, the first obstruction class of the foliation induced on Z by a b -symplectic structure on (M, Z) vanishes.

7.2. The defining two-form of a foliation and the second obstruction class. Now assume that Z is endowed with a regular corank one Poisson structure Π and that \mathcal{F} is the corresponding foliation of Z by symplectic leaves. Furthermore assume that $c_{\mathcal{F}} = 0$ and fix a closed defining one-form α .

Definition 40. A form $\omega \in \Omega^2(Z)$ is a **defining two-form** of the foliation \mathcal{F} induced by the Poisson structure Π if $i_L^* \omega$ is the symplectic form induced by Π on each leaf $L \xrightarrow{i_L} Z$.

When the Poisson structure Π on Z is induced by a b -symplectic structure on (M, Z) , a defining two-form can be chosen such that $\iota_{v_{\text{mod}}|_Z} \omega = 0$.

Proposition 41. *If (M, Z) is a b -symplectic manifold then a defining two-form ω of the symplectic foliation \mathcal{F} on Z that satisfies $\iota_{v_{\text{mod}}|_Z} \omega = 0$ is necessarily closed.*

Proof. The flow of v_{mod} preserves the Poisson structure Π , so it also preserves the corresponding defining two-form ω : $\mathcal{L}_{v_{\text{mod}}|_Z} \omega = 0$. Applying Cartan's formula we have

$$0 = \mathcal{L}_{v_{\text{mod}}|_Z} \omega = d\iota_{v_{\text{mod}}|_Z} \omega + \iota_{v_{\text{mod}}|_Z} d\omega.$$

The first summand vanishes by hypothesis, and so the second summand vanishes as well: $\iota_{v_{\text{mod}}|_Z} d\omega = 0$.

Since for $p \in Z$ and L the corank 2 symplectic leaf through p we have the decomposition

$$T_p Z = \text{span}(v_{\text{mod}}|_Z) \oplus T_p L,$$

it only remains to check that $d\omega(v_1, v_2, v_3) = 0$ for $v_1, v_2, v_3 \in T_p L$. But $i_L^* \omega$ is a symplectic form on L , hence closed, and therefore $d\omega(v_1, v_2, v_3) = 0$. \square

Considering a different volume form on M changes the modular vector field from v_{mod} to $v'_{\text{mod}} = v_{\text{mod}} + u_g$, where u_g is the hamiltonian vector field of the function g , and the defining two-form from ω to $\omega' = \omega + dg \wedge \alpha = \omega + d(g \wedge \alpha)$, thus not changing the cohomology class $[\omega'] = [\omega] \in H^2(Z)$.

We follow [GMP] for the definition of the second obstruction class, an invariant related to the defining two-form of the symplectic foliation \mathcal{F} induced by a regular corank one Poisson structure Π on Z . Because $i_L^* d\omega = d(i_L^* \omega) = 0$ for all $L \in \mathcal{F}$ and α is a defining one-form of the foliation \mathcal{F} , we have

$$d\omega = \mu \wedge \alpha \text{ for some } \mu \in \Omega^2(Z).$$

Furthermore, because $0 = d(d\omega) = d\mu \wedge \alpha - \mu \wedge d\alpha$ and α is closed, the 3-form $d\mu$ is in the subcomplex $\alpha \wedge \Omega(Z)$, and so $d(j\mu) = 0$.

Definition 42. The **second obstruction class** of the foliation \mathcal{F} is

$$\sigma_{\mathcal{F}} = [j\mu] \in H^2(\Omega(Z)/\alpha \wedge \Omega(Z)).$$

One can prove that the class $\sigma_{\mathcal{F}}$ is independent of the choice of ω .

Theorem 43. [GMP] *The second obstruction class vanishes, $\sigma_{\mathcal{F}} = 0$, if and only if one can choose the defining two-form ω of \mathcal{F} to be closed.*

In particular, the second obstruction class (and also the first) of the foliation induced on Z by a b -symplectic structure on (M, Z) vanishes.

This second invariant has also been studied by Gotay [Ga] in the setting of coisotropic embeddings and has a nice interpretation for symplectic fiber bundles (see [GLSW]).

Finally, we recall a result from [GMP] that gives a topological description of a compact Poisson manifold whose regular corank one symplectic foliation contains a compact leaf and which has vanishing first and second invariants. Indeed, in that paper we proved that if the invariants $c_{\mathcal{F}}$ and $\sigma_{\mathcal{F}}$ vanish, then the manifold was the exceptional hypersurface of a Poisson manifold. In this section we have proved the converse, thus establishing the following theorem:

Theorem 44. *If Z is an oriented compact connected regular Poisson manifold of corank one and \mathcal{F} is its symplectic foliation, then $c_{\mathcal{F}} = \sigma_{\mathcal{F}} = 0$ if and only if there exists a Poisson vector field transversal to \mathcal{F} . If furthermore \mathcal{F} contains a compact leaf L , then every leaf of \mathcal{F} is symplectomorphic to L , and Z is the total space of a fibration over \mathbb{S}^1 and the mapping torus⁷ of the symplectomorphism $\phi : L \rightarrow L$ given by the holonomy map of the fibration over \mathbb{S}^1 .*

8. THE EXTENSION PROBLEM

A b -symplectic structure on (M, Z) induces on Z a regular corank one Poisson structure and corresponding foliation by symplectic leaves. Conversely, we can ask: when is a manifold Z , endowed with a regular corank one Poisson structure Π and corresponding symplectic foliation \mathcal{F} , the exceptional hypersurface of a b -symplectic manifold (M, Z) ? And to what extent is such an extension unique? We will see that an extension of Z to a b -symplectic tubular neighbourhood exists exactly when the obstruction classes $c_{\mathcal{F}}$ and $\sigma_{\mathcal{F}}$ of Z vanish, and that uniqueness is related to the modular vector class.

Theorem 45. *Let Π be a regular corank one Poisson structure on a compact manifold Z , and \mathcal{F} the induced foliation by symplectic leaves.*

Then $c_{\mathcal{F}} = \sigma_{\mathcal{F}} = 0$ if and only if Z is the exceptional hypersurface of a b -symplectic manifold (M, Z) whose b -symplectic form induces on Z the Poisson structure Π .

Furthermore, two such extensions (M_0, Z) and (M_1, Z) are b -symplectomorphic on a tubular neighbourhood of Z if and only if the image of their modular vector class under the map below is the same:

$$H_{\text{Poisson}}^1(M) \rightarrow H_{\text{Poisson}}^1(Z).$$

⁷The mapping torus of $\phi : L \rightarrow L$ is the space $\frac{L \times [0,1]}{(x,0) \sim (\phi(x),1)}$.

Proof. We have seen in Section 7 that if Z is the exceptional hypersurface of a b -symplectic manifold (M, Z) , then the obstructions $c_{\mathcal{F}}$ and $\sigma_{\mathcal{F}}$ vanish.

Conversely, assume that $c_{\mathcal{F}} = \sigma_{\mathcal{F}} = 0$ and let α and ω be closed defining one- and two-forms of the foliation \mathcal{F} on Z . Let $M = Z \times (-\varepsilon, \varepsilon)$ and $p : M \rightarrow Z$ be the projection onto the first factor. Define, with $t \in (-\varepsilon, \varepsilon)$,

$$\tilde{\omega} = p^* \alpha \wedge \frac{dt}{t} + p^* \omega.$$

By construction, $\tilde{\omega}$ is a non-degenerate b -form which induces on $Z \times \{0\}$ the given Poisson structure Π . Furthermore, it is closed because α and ω are closed, and so $\tilde{\omega}$ is a b -symplectic form.

Now, let (M_0, Z) and (M_1, Z) be two such extensions, with b -symplectic forms $\tilde{\omega}_0$ and $\tilde{\omega}_1$ respectively. By choosing small enough tubular neighbourhoods of Z in M_0 and M_1 we can assume that we have two b -symplectic structures $\tilde{\omega}_0$ and $\tilde{\omega}_1$ on the same b -manifold (U, Z) . By hypothesis, these induce on Z the same Poisson structure Π , and their corresponding modular vector fields $v_{\text{mod } 0}$ and $v_{\text{mod } 1}$ (for some choice of volume form) are such that $v_{\text{mod } 0}|_Z$ differs from $v_{\text{mod } 1}|_Z$ by a hamiltonian vector field on Z . Then, by Theorem 31, the two structures $\tilde{\omega}_0$ and $\tilde{\omega}_1$ are b -symplectomorphic in a possibly smaller tubular neighbourhood of Z . \square

Example 46. Let $Z = \mathbb{S}^3$ and \mathcal{F} be a codimension-one foliation (for example the Reeb foliation). If the first obstruction class $c_{\mathcal{F}}$ were to vanish, there would exist a closed defining one-form α . But a closed one-form on \mathbb{S}^3 is necessarily exact, $\alpha = df$, and since \mathbb{S}^3 is compact α would vanish at the singular points of f . Thus, \mathbb{S}^3 cannot be the exceptional hypersurface of a b -symplectic manifold.

Example 47. Consider $Z = \mathbb{T}^3$ with coordinates $\theta_1, \theta_2, \theta_3$ and \mathcal{F} the codimension-one foliation on Z with leaves the different k -levels, $k \in \mathbb{R}$, of

$$\theta_3 = a\theta_1 + b\theta_2 + k,$$

where $a, b \in \mathbb{R}$ are fixed and independent over \mathbb{Q} ; then each leaf is diffeomorphic to \mathbb{R}^2 [Ma]. The one-form

$$\alpha = \frac{a}{a^2 + b^2 + 1} d\theta_1 + \frac{b}{a^2 + b^2 + 1} d\theta_2 - \frac{1}{a^2 + b^2 + 1} d\theta_3$$

is a defining one-form for \mathcal{F} and there is a Poisson structure Π on Z which induces the foliation \mathcal{F} and for which

$$\omega = d\theta_1 \wedge d\theta_2 + b d\theta_1 \wedge d\theta_3 - a d\theta_2 \wedge d\theta_3$$

is the defining two-form. Both α and ω are closed, and so the invariants $c_{\mathcal{F}}$ and $\sigma_{\mathcal{F}}$ vanish.

The manifold Z is the exceptional hypersurface of a b -symplectic tubular neighbourhood (\mathcal{U}, Z) with b -symplectic form

$$\tilde{\omega} = \frac{dt}{t} \wedge p^* \alpha + p^* \omega,$$

where $p : \mathcal{U} = Z \times (-\varepsilon, \varepsilon) \rightarrow Z$ is projection and $t \in (-\varepsilon, \varepsilon)$. On Z , the b -symplectic form $\tilde{\omega}$ induces the Poisson structure Π .

To produce a global example of a compact b -symplectic extension of Z , we use instead $Z = \mathbb{T}^3 \times \{0, \pi\}$. Then (\mathbb{T}^4, Z) with

$$\tilde{\omega} = \frac{1}{\sin \theta_4} d\theta_4 \wedge p^* \alpha + p^* \omega$$

is a compact b -symplectic manifold.

9. INTEGRABILITY OF b -SYMPLECTIC MANIFOLDS AS POISSON MANIFOLDS

We begin reviewing the notion of symplectic groupoid as defined by Weinstein in [We87]. We recall that a groupoid is a set Γ equipped with a subset Γ_0 of identity elements, a source map $\alpha : \Gamma \rightarrow \Gamma_0$, a target map $\beta : \Gamma \rightarrow \Gamma_0$, a multiplication operation $(x, y) \rightarrow x \cdot y$ on the set of (x, y) 's satisfying $\beta(x) = \alpha(y)$, and an inversion operation $\iota : \Gamma \rightarrow \Gamma$, with the data above satisfying the obvious generalizations of the usual group axioms.

If Γ is a C^∞ manifold and the data above is also C^∞ , then Γ is called a differentiable groupoid. If Γ is also equipped with a symplectic form Ω for which the manifold $\Gamma_3 = \{(z, x, y) : z = x \cdot y\}$ is Lagrangian in $(\Gamma, \Omega) \times (\Gamma, -\Omega) \times (\Gamma, -\Omega)$ then Γ is called a symplectic groupoid.

It is easy to see from the definition above that Γ_0 is a Lagrangian submanifold of Γ and with a little more effort one can show that Γ_0 has an intrinsic Poisson structure for which the maps α and β are Poisson maps. Thus for every symplectic groupoid Γ one gets an intrinsically associated Poisson manifold Γ_0 , and as pointed out in [We87] it is natural to ask if the converse is true: Given a Poisson manifold Γ_0 can it be “integrated” to a symplectic groupoid Γ having Γ_0 as its set of identity elements? Counterexamples show that this isn't always the case, but that for many familiar examples of Poisson manifolds it holds true. Two examples, which will figure in our application of this theory to b -symplectic manifolds, are the following:

- (1) If $\Gamma_0 = M$ with (M, ω) a symplectic manifold, then one can take Γ to be the pair groupoid $M \times M^-$ with the composition law $(x, y) \cdot (y, z) = (x, z)$.
- (2) Let G be a Lie group, \mathfrak{g} its Lie algebra and $\Gamma_0 = \mathfrak{g}^*$ with its Poisson bivector field

$$f \in \mathfrak{g}^* \rightarrow \Pi_f \in \Lambda^2(T_f \mathfrak{g}^*) \cong \Lambda^2(\mathfrak{g})^*,$$

where $\Pi_f(x \wedge y) = \langle f, [x, y] \rangle$. In this case, the symplectic groupoid integrating Γ_0 is T^*G with source and target maps

$$\alpha, \beta : T^*G \rightarrow G \times \mathfrak{g}^* \rightarrow \mathfrak{g}^*$$

given by the right and left trivializations on T^*G .

In the 1990's, the question posed by Weinstein was formulated in a more general context – “Given a Lie algebroid, can it be integrated to a Lie groupoid?” – and in 2003, necessary and sufficient conditions for this to be true were given by Crainic and Fernandes in [CF03]. In particular, they showed that a Poisson manifold M can be integrated to a symplectic groupoid if its Poisson bivector field

$$m \in M \rightarrow \Pi_m \in \Lambda^2(T_m M)$$

is non-degenerate on an open dense subset of M . Because a b -Poisson bivector field Π is non-degenerate on $M \setminus Z$, all b -symplectic manifolds are integrable.

Nonetheless, it would be nice to have a concrete description of the Γ corresponding to the Γ_0 given by a b -symplectic manifold (M, ω) . Locally such a description can be given in terms of the two examples described earlier due to the fact that the b -symplectic form can be written in a neighbourhood of a point $p \in Z$ as (see Remark 20)

$$\omega = \omega_{L_p} + (\Pi^T)^\sharp.$$

Proposition 48. *The local symplectic groupoid integrating M in a neighbourhood of a point $p \in Z$ is the product groupoid*

$$T^*G \times \Gamma,$$

where G is the “ $ax + b$ group” (with Lie algebra structure given by $[e_1, e_2] = e_2$) and Γ is the pair groupoid $(L_p, \omega_{L_p}) \times (L_p, \omega_{L_p}^-)$.

A semiglobal integrability result, i.e., a description of the symplectic groupoid integrating M in a neighbourhood the exceptional hypersurface Z , rather than the neighbourhood of a point in Z , requires first that we discuss the integrability of Z . This was done in [GMP], and we recall here the main results as applied to the case in study:

Proposition 49. [GMP] *If (M, Z) is a b -symplectic manifold, then the induced regular Poisson manifold Z is integrable.*

This was obtained using [CF04, Corollary 14] and as a consequence of the existence of a leafwise symplectic embedding for Z . This is guaranteed by the vanishing of the second invariant $\sigma_{\mathcal{F}}$, which, as recalled in section 7, can be interpreted via Gotay’s embedding theorem exactly as measuring the obstruction of the existence of a closed two-form on Z which restricts to a symplectic form on each leaf of the foliation.

Theorem 50. [GMP] *If (M, Z) is a b -symplectic manifold and the induced symplectic foliation of Z contains a compact leaf, then all leaves are compact and the hypersurface Z is a mapping torus, and furthermore Z is integrable and its Weinstein's symplectic groupoid is the associated mapping torus groupoid.*

We describe succinctly how the Weinstein's symplectic groupoid for Z is constructed in [GMP]. For a point $p \in Z$ contained in the symplectic leaf L_p , consider $\Sigma = ((L_p \times L_p^-) \times T^*(\mathbb{R}), \Omega + d\lambda_{\text{liouville}})$ with the groupoid product structure. Then, consider a Poisson vector field on Z which generates the \mathbb{S}^1 -action on the mapping torus Z , in our case we may consider $v_{\text{mod}}|_Z$, and use a lift of this vector field to construct the associated mapping torus groupoid. This will give us the Weinstein's symplectic groupoid of Z .

A semiglobal integrability result for b -symplectic manifolds can be obtained by combining Theorem 50 with the semiglobal model for b -symplectic manifolds given by Theorem 44. For this we consider the projection map $p : \mathcal{U} \rightarrow Z$ from a tubular neighbourhood \mathcal{U} of Z and use it to pullback the groupoid structure on Z to \mathcal{U} , which is fairly simple when Z is a mapping torus, as described above.

The description and classification of symplectic groupoids of various Poisson manifolds and in particular of b -symplectic manifolds is recent work of Gualtieri and Li. We refer the reader to their preprint [GL] for further details.

10. INTEGRABLE SYSTEMS ON b -SYMPLECTIC MANIFOLDS

This paper is the second of three papers on b -symplectic geometry. In the first paper we studied the geometry of regular Poisson manifolds with codimension-one symplectic foliations, and in particular proved the results about such manifolds that are quoted in section 7. In the third paper in this series we will study integrable systems and Hamiltonian actions on b -symplectic manifolds and in particular prove the following result: In the case of Poisson b -manifolds, an integrable system is always splitted (in the sense of Weinstein and the integrable system) in a neighbourhood of Z . We then have,

Theorem 51. *An integrable system on a b -symplectic manifold of dimension $2n$ is equivalent to an integrable system with functions (f_1, \dots, f_n) where in a neighbourhood of the critical set Z :*

- (1) *The function f_1 can be chosen to be a defining function for Z .*
- (2) *The remaining first integrals (f_2, \dots, f_n) are functions on Z defining an integrable system on Z with respect to the restricted Poisson structure Π_Z .*

One implication of this result is that most of the classical normal form and action-angle results for symplectic manifolds also hold for b -symplectic manifolds in a neighbourhood of points in Z including possible non-degenerate singularities as studied by Eliasson, Zung and the second author of this paper ([E84, E90, Mi, MZ]). In particular as we will see in the following paper,

Theorem 52. *Given an integrable system with non-degenerate singularities on a b -symplectic manifold (M, Z) , there exist Eliasson-type normal forms in a neighbourhood of points in Z and the minimal rank for these singularities is one along Z .*

REFERENCES

- [Ca] A. Cannas da Silva, *Lectures on Symplectic Geometry* (book), Lecture Notes in Mathematics 1764, Corrected 2nd Printing, Springer, 2008.
- [CW] A. Cannas da Silva and A. Weinstein, *Geometric Models for Noncommutative Algebras* (book), Berkeley Mathematics Lecture Notes series, AMS, 1999.
- [CDW] A. Coste, P. Dazord and A. Weinstein, *Groupoides symplectiques*, Publ. Dept. Math. Univ. Claude-Bernard Lyon I, 1987.
- [CF03] M. Crainic and R.L. Fernandes, *Integrability of Lie brackets*, Annals of Mathematics 157, 2003, pp. 575–620.
- [CF04] M. Crainic and R.L. Fernandes, *Integrability of Poisson brackets*, J. of Differential Geometry 66, 2004, pp. 71–137.
- [CF11] M. Crainic and R.L. Fernandes, *Lectures on Integrability of Lie Brackets*, Geometry and Topology Monographs 17, 2011, pp. 1–107.
- [Co] J. Conn, *Normal forms for smooth Poisson structures*, Annals of Mathematics (2) 121, no. 3, 1985, pp. 565–593.
- [E84] L.H. Eliasson, *Normal forms for Hamiltonian systems with Poisson commuting integrals*, Ph.D. thesis, 1984.
- [E90] L.H. Eliasson, *Normal forms for Hamiltonian systems with Poisson commuting integrals – elliptic case*, Comment. Math. Helv. 65, no. 1, 1990, pp. 4–35.
- [F] P. Freijlich, *h -principles around Poisson geometry*, Ph.D. thesis, Instituto Superior Tecnico, 2011.
- [Ga] M. Gotay, *On coisotropic embeddings of presymplectic manifolds*, Proc. Amer. Math. Soc. 84, 1982, pp. 111–114.
- [Go] R. Goto, *Rozansky-Witten invariants of log symplectic manifolds, integrable systems, topology, and physics*, Contemp. Math., vol. 309, Amer. Math. Soc., Providence RI, 2002, pp. 69–84.
- [GLSW] M. Gotay, R. Lashof, J. Sniatcky, A. Weinstein, *Closed forms on symplectic fibre bundles*, Comment. Math. Helvetici 58, 1983, pp. 617–621.
- [GL] M. Gualtieri and S. Li *Symplectic groupoids of log symplectic manifolds*, preprint 2012.
- [GMP] V. Guillemin, E. Miranda and A.R. Pires, *Codimension one symplectic foliations and regular Poisson structures*, Bulletin of the Brazilian Mathematical Society, New Series 42 (4), 2011, pp. 1–17.
- [K] Y. Kosmann-Schwarzbach, *Poisson manifolds, Lie algebroids, modular classes: a survey*, SIGMA, 4, 005, 2008, 30 pages.
- [Ma] V. Mamaev, *Codimension one foliations of flat 3-manifolds*, Sbornik: Mathematics 187:6, 1996, pp. 823–833.

- [Mt] D. Martinez-Torres, *Global classification of generic multi-vector fields of top degree*, Journal of the London Mathematical Society, 69:3, 2004, pp. 751–766.
- [Me] R. Melrose, *Atiyah-Patodi-Singer Index Theorem* (book), Research Notices in Mathematics, A.K. Peters, Wellesley, 1993.
- [Mi] E. Miranda, *On symplectic linearization of singular Lagrangian foliations*, Ph.D. thesis, Universitat de Barcelona, ISBN: 9788469412374, 2003.
- [MZ] E. Miranda and N.T. Zung, *Equivariant normal forms for nondegenerate singular orbits of integrable Hamiltonian systems*, Ann. Sci. Ecole Norm. Sup., 37, no. 6, 2004, pp. 819–839.
- [NT] R. Nest and B. Tsygan, *Formal deformations of symplectic manifolds with boundary*, J. Reine Angew. Math. 481, 1996, pp. 27–54.
- [R] O. Radko, *A classification of topologically stable Poisson structures on a compact oriented surface*, Journal of Symplectic Geometry, 1, no. 3, 2002, pp. 523–542.
- [V90] I. Vaisman, *Remarks on the Lichnerowicz-Poisson cohomology*, Ann. Inst. Fourier, Grenoble, 40, no. 4, 1990, pp. 951–963.
- [V94] I. Vaisman, *Lectures on the geometry of Poisson manifolds* (book), Progress in Mathematics, 118, Birkhauser Verlag, Basel, 1994.
- [We83] A. Weinstein, *The local structure of Poisson manifolds.*, Journal of Differential Geometry 18, no. 3, 1983, pp. 523–557.
- [We87] A. Weinstein, *Symplectic groupoids and Poisson manifolds*, Bull. Amer. Math. Soc. (N.S.) Volume 16, no. 1, 1987, pp. 101–104.
- [We97] A. Weinstein, *The modular automorphism group of a Poisson manifold*, Journal of Geometry and Physics (23), nos. 3-4, 1997, pp. 379–394.
- [X] P. Xu, *Poisson cohomology of regular Poisson manifolds*, Ann. Inst. Fourier, Grenoble, 42, no. 4, 1992, pp. 967–988.

VICTOR GUILLEMIN, DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE MA, US, *e-mail*: vwg@math.mit.edu

EVA MIRANDA, DEPARTAMENT DE MATEMÀTICA APLICADA I, UNIVERSITAT POLITÈCNICA DE CATALUNYA, BARCELONA, SPAIN, *e-mail*: eva.miranda@upc.edu

ANA RITA PIRES, DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY *e-mail*: apires@math.cornell.edu